

Supersymmetric Unification Without Proton Decay*

G. Lazarides and C. Panagiotakopoulos

Physics Division
School of Technology
University of Thessaloniki
Thessaloniki, Greece

Q. Shafi

Bartol Research Institute
University of Delaware
Newark, Delaware 19716 U.S.A.

Abstract

We present a supersymmetric grand unification scheme based on the gauge group $SU(3)_c \times SU(3)_L \times SU(3)_R$ in which the proton and the lightest supersymmetric particle are stable, neutrinos are necessarily massive, and the observed baryon asymmetry originates in the lepton sector. Such models are also consistent with the measured value of $\sin^2 \theta_W$ as well as unification of the gauge couplings.

*Supported in part by NATO Grant #CRG-910573 and Department of Energy Grant #DE-FG02-91ER406267.

The minimal supersymmetric $SU(5)$ model⁽¹⁾ predicts a value for $\sin^2 \theta_W$ which is remarkably consistent with the measured numbers⁽²⁾. Despite this success, however, it appears that the minimal scheme faces an uphill task. For instance, the predicted asymptotic mass relations involving the first two families ($m_d = m_e$ and $m_s = m_\mu$) are in glaring contradiction with the observations. Furthermore, there is a close relationship between the light fermion masses and proton decay (into $K^+ \bar{\nu}_i$, etc.) mediated through dimension five interactions. The experimental situation regarding the latter is getting to a point⁽³⁾ where it appears puzzling why the spectacular events have not been observed. Additional motivation for thinking beyond the minimal $SU(5)$ framework comes from considerations of neutrino masses, baryon asymmetry, etc.

One modest extension would be an enlargement of the Higgs sector as well as the introduction of singlet chiral superfields. Another possibility is to embed in an $SO(10)$ framework. In this paper we follow a relatively less treaded path, in which the standard $SU(3)_c \times SU(2)_L \times U(1)_Y$ model is embedded in $G \equiv SU(3)_c \times SU(3)_L \times SU(3)_R$. There are additional reasons besides those already listed for studying models based on G . The simplest compactification schemes⁽⁴⁾ involving the heterotic $E_8 \times E_8$ superstring theory give rise to G , although the present study will be largely motivated within the ordinary GUT framework. The imposition of a Z_2 symmetry ensures the (almost complete) absence of baryon number violation. Indeed, the main source of baryon number violation is the non-perturbative sector of the standard electroweak model! Other ‘advantages’ include the absence of wrong fermion mass relations, ‘naturally’ massive neutrinos, and an elegant mechanism for

generating the observed baryon asymmetry. We will present here two different realizations of the $G \equiv SU(3)_c \times SU(3)_L \times SU(3)_R$ model. In the first one the symmetry breaking of G to the minimal supersymmetric standard model (MSSM) proceeds in two steps, whereas in the second scenario there is no intermediate step breaking.

The left handed lepton, quark and antiquark superfields transform under G as follows:

$$\begin{aligned}\lambda &= (1, \bar{3}, 3) = \begin{pmatrix} H^{(1)} & H^{(2)} & L \\ E^c & \nu^c & N \end{pmatrix}, \\ Q &= (3, 3, 1) = \begin{pmatrix} q \\ g \end{pmatrix}, \\ Q^c &= (\bar{3}, 1, \bar{3}) = \begin{pmatrix} u^c & d^c & g^c \end{pmatrix}.\end{aligned}\tag{1}$$

Here N and ν^c denote standard model singlet superfields, while g (g^c) is an additional down-type quark (antiquark). We will be working with five λ 's, two $\bar{\lambda}$'s, five Q 's, two $\bar{Q}$'s, five Q^c 's, two $\bar{Q}^c$'s and two G -singlet superfields S_1, S_2 . Notice that the field content is chosen to ensure unification above the GUT scale.

We choose to impose invariance under the discrete Z_2 symmetry⁽⁵⁾ P_1 generated by $Q_a \rightarrow -Q_a$, $Q_a^c \rightarrow -Q_a^c$, $\lambda_a \rightarrow \lambda_a$ ($a = 1, 2, \dots, 5$), $\bar{Q}_\alpha \rightarrow -\bar{Q}_\alpha$, $\bar{Q}_\alpha^c \rightarrow -\bar{Q}_\alpha^c$, $\bar{\lambda}_\alpha \rightarrow \bar{\lambda}_\alpha$ ($\alpha = 1, 2$). This has the effect of forbidding baryon number violating couplings $Q_{a_1} Q_{a_2} Q_{a_3}$, $Q_{a_1}^c Q_{a_2}^c Q_{a_3}^c$, $\bar{Q}_{\alpha_1} \bar{Q}_{\alpha_2} \bar{Q}_{\alpha_3}$ and $\bar{Q}_{\alpha_1}^c \bar{Q}_{\alpha_2}^c \bar{Q}_{\alpha_3}^c$. Consequently, proton decay mediated by dimension four, five and six operators cannot occur. Moreover, the structure of G is such that there is no gauge boson mediated decay. The proton is stable.

The symmetry breaking to MSSM is obtained through superpotential

couplings of the form $S_1(\lambda_4\bar{\lambda}_1 - M_X^2)$ and $S_2(\lambda_5\bar{\lambda}_2 - M_{B-L}^2)$. The superlarge vacuum expectation values are acquired by the following fields:

$$\begin{aligned} \langle N_4 \rangle &= \langle \bar{N}_1 \rangle^* = M_X, \\ \langle \nu_5^c \rangle &= \langle \bar{\nu}_2^c \rangle^* = M_{B-L}. \end{aligned} \quad (2)$$

We also introduce in the superpotential explicit mass terms of order M_X for $Q, \bar{Q}$ and $Q^c, \bar{Q}^c$ pairs. In the first realization of the theory, we assume that $M_{B-L} \ll M_X$. This means that the breaking to MSSM proceeds in two steps:

$$G \xrightarrow{M_X} SU(3)_c \times SU(2)_L \times SU(2)_R \times U(1)_{B-L} \xrightarrow{M_{B-L}} MSSM. \quad (3)$$

We will shortly provide estimates for the scales M_X and M_{B-L} .

In addition to the Z_2 symmetry P_1 , we impose an additional Z_2 symmetry P_2 , under which $\lambda_5 \rightarrow -\lambda_5$, $\bar{\lambda}_2 \rightarrow -\bar{\lambda}_2$, while all other fields remain invariant. This symmetry, combined with the Z_2 subgroup generated by the element $(\text{diag}(-1, -1), \text{diag}(-1, -1))$ of $SU(2)_L \times SU(2)_R$, remains unbroken by all the vacuum expectation values and acts as ‘matter parity’. The purpose of introducing this symmetry is to eliminate some lepton number violating couplings at the level of the MSSM. This is necessary for the implementation of our mechanism of generating the baryon asymmetry of the Universe. The physical consequences of matter parity will become clear in the following.

We next study the spectrum of the theory for the case $M_{B-L} \ll M_X$. The mass matrices will be classified according to the quantum numbers of the states under the gauge group $SU(3)_c \times SU(2)_L \times U(1)_Y$ and the Z_2 matter parity. We will also confine ourselves to the discussion of the fermionic mass

matrices. The result applies to the bosonic components as well, modulo corrections of the order of the supersymmetry breaking scale $M_S \sim 1TeV$.

Let us first look at the mass matrix of the colorless $SU(2)_L$ - doublet fields with positive matter parity. The bosonic partners of these fields have the correct quantum numbers to play the role of the electroweak Higgs doublets. The mass matrix couples $H_m^{(2)}$ ($m = 1, 2, 3, 4$), $\bar{H}_1^{(1)}$ and L_5 with $H_m^{(1)}$ ($m = 1, 2, 3, 4$), $\bar{H}_1^{(2)e}$ and $\bar{L}_2$. In the limit $M_{B-L} = 0$, L_5 and $\bar{L}_2$ are exactly massless and decouple from the rest of the mass matrix. The remaining mass matrix is symmetric with all eigenvalues typically $\sim M_X$. Mixings between $H_m^{(2)}$ and $\bar{H}_1^{(1)}$, as well as between $H_m^{(1)}$ and $\bar{H}_1^{(2)}$, are typically $\sim M_X/M_P$ ($M_P \simeq 1.2 \cdot 10^{19} GeV$ is the Planck mass), whereas mixings among the $H_m^{(2)}$'s and among the $H_m^{(1)}$'s are of order unity. Diagonalization of this symmetric mass matrix by a unitary transformation gives a positive matrix, $\text{diag} (M_1, M_2, M_3, M_4, \bar{M}_1)$, which predominantly couples four linear combinations of the four $H_m^{(2)}$'s with the corresponding four linear combinations of the four $H_m^{(1)}$'s as well as $\bar{H}_1^{(1)}$ with $\bar{H}_1^{(2)}$. To obtain a unique light pair of electroweak Higgs doublets, $h^{(2)}$, $h^{(1)}$, we must fine tune one eigenvalue among the four M 's so that it becomes $\sim M_S$. Thus we reach the important conclusion that, neglecting mass contributions which vanish in the $M_{B-L} = 0$ limit, the pair of electroweak doublets, $h^{(2)}$, $h^{(1)}$, predominantly lies in one linear combination λ_0 of λ 's. Removing the superheavy states with masses of order M_X , we are left with a 2×2 mass matrix which couples $h^{(2)}$ and L_5 with $h^{(1)}$ and $\bar{L}_2$. As M_{B-L} grows, the $L_5 - h^{(1)}$, $h^{(2)} - \bar{L}_2$ and $L_5 - \bar{L}_2$ mass terms become of the order of M_{B-L} , $M_X M_{B-L}/M_P$ and M_{B-L}^2/M_P respectively. In order to keep the pair of electroweak doublets, $h^{(2)}$, $h^{(1)}$, essentially

unaffected by these contributions, we must then fine tune the off-diagonal mass terms so that both of them are much smaller than $\sim M_{B-L}^2/M_P$ and their product is lower than $\sim M_S M_{B-L}^2/M_P$.

The mass matrix of the colorless $SU(2)_L$ - doublets with negative matter parity couples $H_5^{(2)}$, $\bar{H}_2^{(1)}$ and L_m ($m = 1, 2, 3, 4$) with $H_5^{(2)}$, $\bar{H}_2^{(2)}$ and $\bar{L}_1$. The fields $H_5^{(2)}$ and $\bar{H}_2^{(1)}$ predominantly pair up with $H_5^{(1)}$ and $\bar{H}_2^{(2)}$ to form states with masses $\sim M_X$. Also, one combination of the four L_m 's pairs up with $\bar{L}_1$ to form a state with mass $\sim M_X^2/M_P$. The remaining three orthogonal linear combinations l_i ($i = 1, 2, 3$) of the four L_m 's remain massless and play the role of the ordinary light lepton doublets.

The $SU(2)_L$ - singlet leptons with positive matter parity are E_5^c and $\bar{E}_2^c$. They form a state with mass $\sim M_{B-L}^2/M_P$. One linear combination of the four negative matter parity singlet leptons E_m^c ($m = 1, 2, 3, 4$) pairs up with $\bar{E}_1^c$ to form a state with mass $\sim M_X^2/M_P$. The three remaining orthogonal linear combinations e_i^c ($i = 1, 2, 3$) which belong, to a very good approximation, to the same λ 's as the light l_i ($i = 1, 2, 3$), remain massless and play the role of the ordinary light $SU(2)_L$ -singlet leptons.

In the positive matter parity neutral sector (N_m ($m = 1, 2, 3, 4$), $\bar{N}_1$, ν_5^c , $\bar{\nu}_2^c$), all states get masses $\sim M_X^2/M_P$ except for the two linear combinations acquiring the large expectation values which get masses of the order of the corresponding expectation value (M_X and M_{B-L} , respectively) and one combination of ν_5^c and $\bar{\nu}_2^c$ which gets a mass $\sim M_{B-L}^2/M_P$. In the negative matter parity neutral sector (N_5 , $\bar{N}_2$, ν_m^c ($m = 1, 2, 3, 4$), $\bar{\nu}_1^c$), N_5 , $\bar{N}_2$ get masses $\sim M_X^2/M_P$ and have mixings $\sim M_{B-L}/M_X$ with ν_m^c 's and $\bar{\nu}_1^c$. One linear combination of the four ν_m^c 's (let us say ν_4^c) pairs up with $\bar{\nu}_1^c$ to form

a state with mass $\sim M_X^2/M_P$. The remaining three linear combinations $\nu_i^c (i = 1, 2, 3)$, which are $SU(2)_R$ - partners of the light e_i^c 's acquire masses $\sim M_{B-L}^2/M_P$. The mass matrix of the light ν_i^c 's is symmetric and therefore can be diagonalized by a single unitary transformation. In the following we will assume that the mass matrix of the ν_i^c 's is already diagonal in the form $\text{diag} (M_{\nu_1^c}, M_{\nu_2^c}, M_{\nu_3^c})$.

The quark sector is simple. After pairing two $Q, \bar{Q}$ and two $Q^c, \bar{Q}^c$ pairs with mass terms assumed of order $10^{-2} M_X$, we are left with three massless doublets $q_i = \begin{pmatrix} u_i \\ d_i \end{pmatrix}$, as well as three massless u_i^c 's and three massless d_i^c 's belonging to the same three Q_i^c 's ($i = 1, 2, 3$), all with negative matter parity. We also have the corresponding three pairs of g_i, g_i^c 's ($i = 1, 2, 3$) which have positive matter parity and acquire 'superheavy' masses $\sim M_X$.

Between the energy scales M_W and M_X , the mass spectrum is not symmetric in the three $SU(3)$'s (see above). However, above M_X the symmetry is restored and the three gauge couplings are equal (here we assume that G emerges at M_P with a unique gauge coupling constant). Using the standard one loop renormalization group evolution of the gauge couplings, and assuming that the supersymmetry breaking scale $M_S \simeq 10^3 \text{GeV}$, one can obtain the following (consistent) solution:

$$\begin{aligned} \sin^2 \theta_W(M_Z) &\approx 0.23, \alpha_s(M_Z) \simeq 0.115, \alpha_G(M_P) \simeq 0.158, \\ M_X &\approx 10^{16.5} \text{GeV}, M_{B-L} \simeq 10^{13.5} \text{GeV}, \end{aligned} \tag{4}$$

where α_G denotes the unified gauge coupling above M_X .

The fact that the pair of electroweak doublets $h^{(2)}, h^{(1)}$ predominantly belongs to the same linear combination of λ_a 's (namely λ_0) together with the

observation that the ‘light’ u_i^c ’s, d_i^c ’s belong to the same three Q_i^c ’s ($i = 1, 2, 3$) implies that the tree-level up- and down-quark mass matrices are proportional (they both come from the same couplings $Q_i Q_j^c \lambda_0$). Assuming that the third generation acquires tree-level masses, we obtain the asymptotic relation

$$\frac{m_t}{m_b} \simeq \tan \beta \equiv \frac{< h^{(1)} >}{< h^{(2)} >}. \quad (5)$$

To account for the correct masses and mixings of the two lighter quark families, we must invoke some other mechanism (radiative corrections, ...) which we will not specify here. In the leptonic sector, we have seen that the three ‘light’ ν_i^c ’s and the light e_i^c ’s belong to the same λ_i ’s ($i = 1, 2, 3$). Thus, the tree-level Dirac mass matrix of the neutrinos is proportional to the tree-level charged lepton mass matrix (since both these mass matrices come from the same couplings $\lambda_i \lambda_j \lambda_0$). Assuming again that only the third family acquires tree-level masses we obtain the asymptotic relation

$$\frac{m_{\nu_\tau}^D}{m_\tau} \simeq \tan \beta \quad (6)$$

where the superscript D denotes a Dirac neutrino mass.

So far we have seen that the model is consistent with the low energy phenomenology. The proton is stable and $\sin^2 \theta_W$ is in the right ball park. Next we address the issue of baryon asymmetry. Since the conventional baryon number violating couplings are missing, we must resort to the lepton sector which contains the lepton number violating Majorana mass terms. The idea then is to generate an initial lepton asymmetry n_L/s through the decays of ν_i^c ($i = 1, 2, 3$) ^(6,7). The non-perturbative sphaleronlike effects in the

electroweak sector⁽⁸⁾ would then convert a fraction of the asymmetry n_L/s into the observed baryon asymmetry n_b/s , provided the ‘reheat’ temperature T_r after the completion of inflation is greater than or equal to about 100 GeV. In order to ensure the survival of n_L/s , we must ensure that all lepton number violating 2-2 scatterings are out of equilibrium at temperatures between T_r and 100 GeV. This is elegantly achieved with the help of the Z_2 matter parity.

Within an inflationary framework driven by a gauge singlet field ϕ (note that inflation is needed here to eliminate the monopole problem), we assume copious production of ν_i^c from the out of equilibrium decay of ϕ as it oscillates about the minimum of the potential. If $m_\phi \gtrsim 2M_{\nu_i^c}$ ($i = 1, 2, 3$), the inflaton will predominantly decay to the heaviest ν_i^c which we assume is ν_3^c (recall that we are in a basis in which the ν_i^c -mass matrix is diagonal). The decay width of ϕ is given by

$$\Gamma_\phi \sim \frac{1}{16\pi} \left(\frac{M_{\nu_3^c}}{\langle \phi \rangle} \right)^2 m_\phi, \quad (7)$$

where $\langle \phi \rangle \sim 10^{17} - 10^{18}$ GeV as estimated from the requirement that $\delta\rho/\rho \sim 10^{-5}$. The corresponding ‘reheat’ temperature is given by

$$\begin{aligned} T_r &\sim \frac{1}{3}(\Gamma_\phi M_P)^{\frac{1}{2}} \\ &\sim 2.6 \times 10^3 \left(\frac{M_{\nu_3^c}}{10^8 \text{ GeV}} \right) \left(\frac{10^{17.5} \text{ GeV}}{\langle \phi \rangle} \right) \left(\frac{m_\phi}{10^{9.5} \text{ GeV}} \right)^{\frac{1}{2}} \text{ GeV}. \end{aligned} \quad (8)$$

We will choose $M_{\nu_3^c} \simeq 10^8$ GeV, $m_\phi \simeq 10^{9.5}$ GeV and $\langle \phi \rangle \simeq 10^{17.5}$ GeV. This implies $T_r \simeq 2.6 \text{ TeV}$, which is higher than the minimum allowed ‘reheat’ temperature.

A simple double-cut diagram which leads to a lepton asymmetry through

ν_i^c decays (in our case $i = 3$) is shown in Fig.1 (see also ref. 6). A straightforward estimate leads to

$$n_{L_i}/s \sim \frac{1}{4\pi} \frac{T_r}{m_\phi} \frac{M_{\nu_i^c}}{\left(\frac{m_D m_D^\dagger}{|\langle h^{(1)} \rangle|^2}\right)_{ii}} \text{Im} \sum_k \left(\frac{m_D m_D^\dagger}{|\langle h^{(1)} \rangle|^2} \right)_{ik}^2 \frac{M_{\nu_k^c}}{\vec{p}^2 + M_{\nu_k^c}^2}, \quad (9)$$

where m_D is the Dirac mass matrix of the neutrinos and the internal three-momentum $|\vec{p}| \simeq \frac{1}{2} M_{\nu_i^c}$. To help maximize the lepton asymmetry we set $M_{\nu_2^c} \simeq 2 \cdot 10^5 \text{ GeV}$, and assume that the dominant matrix element of m_D is $(m_D)_{23}$ (the element which couples ν_2^c and ν_3). Eq. (9) then gives

$$n_{L_3}/s \sim \frac{1}{\pi} \cdot \frac{T_r}{m_\phi} \cdot \frac{M_{\nu_2^c}}{M_{\nu_3^c}} \cdot \left| \frac{(m_D)_{23}}{\langle h^{(1)} \rangle} \right|^2. \quad (10)$$

Proportionality of m_D and the charged lepton mass matrix at tree-level implies

$$\frac{(m_D)_{23}}{\langle h^{(1)} \rangle} \simeq \frac{m_\tau}{\langle h^{(2)} \rangle} \simeq \frac{m_\tau \tan \beta}{\langle h^{(1)} \rangle}. \quad (11)$$

Taking $\tan \beta \simeq 44$ and $\langle h^{(1)} \rangle \simeq 170 \text{ GeV}$, for example, one finds that $(m_D)_{23}/\langle h^{(1)} \rangle \simeq 0.46$ and $n_{L_3}/s \sim 10^{-10} \sim n_b/s$.

With this choice of the parameters, the mass of the tau neutrino turns out to be

$$m_{\nu_\tau} \simeq \frac{(m_\tau \tan \beta)^2}{M_{\nu_2^c}} \sim 30 \text{ MeV}. \quad (12)$$

The requirement of a sizeable baryon asymmetry has led us to a τ -neutrino mass in the MeV range. The lifetime and branching ratio into radiative

modes of such a neutrino is strongly constrained by astrophysical considerations⁽⁹⁾. A neutrino with mass $\sim 30 MeV$ could be astrophysically acceptable if it decays with a lifetime $\tau_{\nu_\tau} \leq 20 sec$ into $e^+e^-\nu$ or $\nu\nu'\bar{\nu}''$, with ν, ν', ν'' any of the two lighter neutrinos. This can easily be achieved provided M_S is somewhat lower than $1TeV$. The relevant diagrams are simple one loop box diagrams involving supersymmetric partners of charged leptons or of neutrinos and two winos.

As another possibility, one can consider the mode $\nu_\tau \rightarrow \nu_e a$, where a is an axion field. The decay rate is then $\Gamma \sim \frac{1}{16\pi}(\frac{m_{\nu_\tau}}{f_a})^2 m_{\nu_\tau}$, where f_a is the axion decay constant. For $f_a \simeq 3 \cdot 10^9 GeV$, we obtain $\tau_{\nu_\tau} \simeq 20 sec$ which is acceptable. Note that an axion may eventually be needed to solve the strong CP problem.

One important consequence of the intermediate scale M_{B-L} is that the ν_i^c masses can be of order $10^8 GeV$, which facilitates leptogenesis to proceed in an inflationary scenario⁽⁷⁾. It also opens up the exciting possibility that one of the ‘light’ neutrinos has mass in the 3-10 eV range, a feature hinted⁽¹⁰⁾ at by the recent COBE data. However, this would mean that the MSW⁽¹¹⁾ solution of the solar neutrino problem cannot be implemented.

The inflationary scenario predicts that the density parameter Ω is unity, and it has been suggested that $\Omega_\nu \approx 0.2 - 0.3$. That is, the bulk of the missing mass is in ‘cold’ dark matter. One of the major consequences of the Z_2 matter parity in our model is the stability of the lightest supersymmetric particle (LSP). The latter can be either the neutralino or possibly the sneutrino. In the first case it has positive matter parity but it has to decay (by fermion number conservation) to an odd number of negative matter

parity light fermions. In the second case it has negative matter parity but has to decay to an even number of negative matter parity light fermions. In both cases the decay process is forbidden by the unbroken Z_2 matter parity. Therefore the LSP is a ‘cold’ dark matter candidate in our model.

In the second realization of the $SU(3)_c \times SU(3)_L \times SU(3)_R$ model, we take $M_{B-L} = M_X$. Assuming that the $Q, \bar{Q}$ and $Q^c, \bar{Q}^c$ mass terms are $\frac{1}{3}M_X$ and the g, g^c ’s acquire masses $\sim 10^{-1}M_X$ one loop renormalization group analysis yields ($M_S \simeq 10^3 GeV$)

$$\begin{aligned} \sin^2 \theta_W(M_Z) &\simeq 0.23, \alpha_s(M_Z) \simeq 0.11, \alpha_G(M_P) \simeq 0.20, \\ M_X &= M_{B-L} \simeq 10^{15} GeV. \end{aligned} \tag{13}$$

In this case, however, the pair of Higgs doublets $h^{(2)}, h^{(1)}$ does not necessarily belong to the same linear combination of λ_a ’s. Also, the light e_i^c ’s and the ‘light’ ν_i^c ’s do not belong to the same three λ_i ’s. This means that we lose the asymptotic relations in eqs. (5) and (6). The ‘heavy’ neutrino masses are $M_{\nu_i^c} \sim M_{B-L}^2/M_P \sim 10^{11} GeV$, and we assume that they are all comparable. The ‘reheat’ temperature is again calculated from formulae (7) and (8), and turns out to be much higher than $1 TeV$. However, taking into account the fact that $M_{\nu^c}/\langle \phi \rangle$ is an upper bound on the effective $\phi \nu_i^c \nu_j^c$ coupling constants, the value of T_r calculated this way should be considered only as an upper bound.

The lepton asymmetry generated by the decaying ν_i^c ’s is given by

$$\frac{n_L}{s} = \sum_{i=1}^3 \frac{n_{Li}}{s}, \tag{14}$$

where the expression for n_{Li}/s is given in eq. (9). All of the ν_i^c ’s have

comparable masses of order 10^{11} GeV , and we can assume that they are equally produced through the decay of the inflaton field ϕ (with mass $m_\phi \gtrsim 2 \cdot 10^{11} \text{ GeV}$). Taking $(m_D)_{33}$ to be the largest element of m_D , the lepton asymmetry, predominantly produced through the decays of $\nu_{1,2}^c$, is estimated to be

$$\frac{n_L}{s} \sim \frac{1}{\pi} \frac{T_r}{m_\phi} \left| \frac{(m_D)_{33}}{\langle h^{(1)} \rangle} \right|^2. \quad (15)$$

The allowed values for $(m_D)_{33}$ satisfy the following inequality

$$m_{\nu_\tau} \simeq \frac{|(m_D)_{33}|^2}{M_{\nu^c}} \lesssim 10 \text{ eV}. \quad (16)$$

Thus, eq.(15) can easily lead to acceptable values for the baryon asymmetry of the Universe. The bound in eq.(16) is saturated for $(m_D)_{33} \simeq 30 \text{ GeV}$, leading to a τ -neutrino mass of about 10 eV which can provide the ‘hot’ dark matter of the universe. The LSP is again a ‘cold’ dark matter candidate.

To conclude, we have outlined a possible alternative to minimal supersymmetric $SU(5)$ unification. It offers the feature that the proton is stable, and it accomplishes this in a relatively elegant manner, avoiding in the process also some unacceptable mass relations. Other advantages include the ease with which the baryon asymmetry is generated (necessarily via leptogenesis) and the presence of some hot dark matter in the form of massive neutrinos. Also, since the Higgs doublets and the color triplet fields belong to different representations, the doublet - triplet splitting can, in principle, be accomplished more easily than in $SU(5)$.

We considered two alternative realizations of the model. In the first

case, the B-L breaking scale is much lower than the unification scale. This leads to asymptotic relations like $m_t/m_b \simeq \tan\beta$. The requirement of a sizable baryon asymmetry in this case favors an unstable τ -neutrino with mass of about 30 MeV and a lifetime of 20 sec. The ν_μ -neutrino mass can be chosen so that it provides either the ‘hot’ dark matter of the universe or is consistent with the MSW solution of the solar neutrino problem. In the second realization of the model the B-L breaking scale coincides with the unification scale. In this case a tau neutrino with mass of order 10eV can provide some ‘hot’ dark matter, while the muon neutrino plays a role in the MSW effect. In either case a stable LSP provides for a ‘cold’ dark matter candidate.

Acknowledgement: We thank K.S. Babu for several important discussions.

References

1. S. Dimopoulos and H. Georgi, Nucl. Phys., B193 (1981) 150; N. Sakai, Z. Phys., C11 (1981) 153.
2. J. Ellis, S. Kelley and D.V. Nanopoulos, Phys. Lett., B249 (1990) 441; V. Amaldi, W. de Boer and H. Furstenan, Phys. Lett., B260 (1991) 447; P. Langacker and M.X. Luo, Phys. Rev. D, D44 (1991) 817.
3. Kamiokande Collaboration (1992).
4. P. Candelas, G. Horowitz, A. Strominger and E. Witten, Nucl. Phys., B258 (1985) 46; E. Witten, Nucl. Phys., B258 (1985) 75.

5. Q. Shafi and X.M. Wang, 1990 (unpublished); L. Ibanez and G.G. Ross, Nucl. Phys., B368 (1992) 3.
6. M. Fukugita and T. Yanagida, Phys. Lett., 174B (1986) 45.
7. G. Lazarides and Q. Shafi, Phys. Lett., 258B (1991) 305.
8. V. Kuzmin, V. Rubakov and M. Shaposhnikov, Phys. Lett., B155 (1985) 36.
9. S.A. Bludman, Phys. Rev., D45 (1992) 4720.
10. R. Schaefer and Q. Shafi, Nature, 359 (1992) 359.
11. L. Wolfenstein, Phys. Rev., D17 (1978) 2369; Phys. Rev., D20 (1979) 2634; S.P. Mikheyev and A.Yu. Smirnov, Usp. Fiz. Nauk., 153 (1987) 3.